

Resource Summary Report

Generated by [RRID](#) on Aug 5, 2026

Parvalbumin antibody

RRID:AB_369891

Type: Antibody

Proper Citation

GeneTex Cat# GTX11427, RRID:AB_369891

Antibody Information

URL: http://antibodyregistry.org/AB_369891

Proper Citation: GeneTex Cat# GTX11427, RRID:AB_369891

Target Antigen: Parvalbumin antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunoprecipitation; Immunohistochemistry - fixed; Western Blot; ELISA; For ELISA: Use at an assay dependent dilution. For IHC-P: Use at a concentration of 1 µg/ml. For IP: Use at a concentration of 5 µg/ml. For WB: Use at a concentration of 0.1 µg/ml. Detects a band of approximately 12 kDa, representing parvalbumin from rat cerebellum. Parvalbumin protein is relatively small and, therefore, it is recommended that electrophoresis be performed using tricine-SDS-PAGE gels and transferred to a nylon membrane. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher. ELISA, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Antibody Name: Parvalbumin antibody

Description: This polyclonal targets Parvalbumin antibody

Target Organism: rat

Antibody ID: AB_369891

Vendor: GeneTex

Catalog Number: GTX11427

Record Creation Time: 20250819T211746+0000

Ratings and Alerts

No rating or validation information has been found for Parvalbumin antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Immunoprecipitation; Immunohistochemistry - fixed; Western Blot; ELISA; For ELISA: Use at an assay dependent dilution. For IHC-P: Use at a concentration of 1 µg/ml. For IP: Use at a concentration of 5 µg/ml. For WB: Use at a concentration of 0.1 µg/ml. Detects a band of approximately 12 kDa, representing parvalbumin from rat cerebellum. Parvalbumin protein is relatively small and, therefore, it is recommended that electrophoresis be performed using tricine-SDS-PAGE gels and transferred to a nylon membrane. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher. ELISA, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.