

Resource Summary Report

Generated by [RRID](#) on Aug 1, 2026

[FLI1 antibody](#)

RRID:AB_370380

Type: Antibody

Proper Citation

GeneTex Cat# GTX15289, RRID:AB_370380

Antibody Information

URL: http://antibodyregistry.org/AB_370380

Proper Citation: GeneTex Cat# GTX15289, RRID:AB_370380

Target Antigen: FLI1 antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunoprecipitation; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; For IHC-P: Use at a dilution of 1:50. Perform enzymatic antigen retrieval before commencing with IHC staining protocol, by boiling tissue sections in 10mM citrate buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min. For IP: Use at a concentration of 2 µg/mg protein lysate. For WB: Use at a concentration of 2 µg/ml for 2hrs. Predicted molecular weight: 54 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Antibody Name: FLI1 antibody

Description: This polyclonal targets FLI1 antibody

Target Organism: rat, mouse, human

Antibody ID: AB_370380

Vendor: GeneTex

Catalog Number: GTX15289

Record Creation Time: 20250820T002032+0000

Record Last Update: 20250820T080053+0000

Ratings and Alerts

No rating or validation information has been found for FLI1 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunoprecipitation; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; For IHC-P: Use at a dilution of 1:50. Perform enzymatic antigen retrieval before commencing with IHC staining protocol, by boiling tissue sections in 10mM citrate buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min. For IP: Use at a concentration of 2 µg/mg protein lysate. For WB: Use at a concentration of 2 µg/ml for 2hrs. Predicted molecular weight: 54 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.