

Resource Summary Report

Generated by [RRID](#) on Aug 2, 2026

VEGF receptor 1 antibody

RRID:AB_370565

Type: Antibody

Proper Citation

GeneTex Cat# GTX22350, RRID:AB_370565

Antibody Information

URL: http://antibodyregistry.org/AB_370565

Proper Citation: GeneTex Cat# GTX22350, RRID:AB_370565

Target Antigen: VEGF receptor 1 antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry - fixed; Immunohistochemistry; Western Blot; IHC: This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining. IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot.

Antibody Name: VEGF receptor 1 antibody

Description: This polyclonal targets VEGF receptor 1 antibody

Target Organism: rat, mouse, human

Antibody ID: AB_370565

Vendor: GeneTex

Catalog Number: GTX22350

Record Creation Time: 20250820T012324+0000

Record Last Update: 20250820T104809+0000

Ratings and Alerts

No rating or validation information has been found for VEGF receptor 1 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry - fixed; Immunohistochemistry; Western Blot; IHC: This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions and antibody dilutions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining. IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.