

Resource Summary Report

Generated by [RRID](#) on Jul 31, 2026

Integrin beta 5 antibody

RRID:AB_371972

Type: Antibody

Proper Citation

GeneTex Cat# GTX15459, RRID:AB_371972

Antibody Information

URL: http://antibodyregistry.org/AB_371972

Proper Citation: GeneTex Cat# GTX15459, RRID:AB_371972

Target Antigen: Integrin beta 5 antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry; Immunoprecipitation; Western Blot; IF: Use at an assay dependent dilution. IHC-P: Use at a dilution of 1/50. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate, pH 6.0 for 10 min followed by cooling at RT for 20 min. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 95.9 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Antibody Name: Integrin beta 5 antibody

Description: This polyclonal targets Integrin beta 5 antibody

Target Organism: rat, hamster, mouse, human

Antibody ID: AB_371972

Vendor: GeneTex

Catalog Number: GTX15459

Record Creation Time: 20250820T001016+0000

Record Last Update: 20250820T073334+0000

Ratings and Alerts

No rating or validation information has been found for Integrin beta 5 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry; Immunoprecipitation; Western Blot; IF: Use at an assay dependent dilution. IHC-P: Use at a dilution of 1/50. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate, pH 6.0 for 10 min followed by cooling at RT for 20 min. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 95.9 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.