

Resource Summary Report

Generated by [RRID](#) on Aug 1, 2026

NGF receptor [NGFR5] antibody

RRID:AB_372631

Type: Antibody

Proper Citation

GeneTex Cat# GTX23125, RRID:AB_372631

Antibody Information

URL: http://antibodyregistry.org/AB_372631

Proper Citation: GeneTex Cat# GTX23125, RRID:AB_372631

Target Antigen: NGF receptor [NGFR5] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: FACS: Use at an assay dependant dilution. IF: Use at an assay dependant dilution. IHC-P: Use at a concentration of 1 - 2 µg/ml. Has not been shown to work in WB. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min., FACS, IF, IHC-P, Flow cytometry/FACS, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined.; Immunohistochemistry; Immunofluorescence; Immunohistochemistry - fixed; Flow Cytometry

Antibody Name: NGF receptor [NGFR5] antibody

Description: This monoclonal targets NGF receptor [NGFR5] antibody

Target Organism: feline, baboon, cat, rabbit, ferret, human

Antibody ID: AB_372631

Vendor: GeneTex

Catalog Number: GTX23125

Record Creation Time: 20250820T051238+0000

Record Last Update: 20250820T210409+0000

Ratings and Alerts

No rating or validation information has been found for NGF receptor [NGFR5] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: FACS: Use at an assay dependant dilution. IF: Use at an assay dependant dilution. IHC-P: Use at a concentration of 1 - 2 µg/ml. Has not been shown to work in WB. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min., FACS, IF, IHC-P, Flow cytometry/FACS, Immunofluorescence, IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined.; Immunohistochemistry; Immunofluorescence; Immunohistochemistry - fixed; Flow Cytometry

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.