

Resource Summary Report

Generated by [RRID](#) on Sep 10, 2026

TGF beta 2 antibody

RRID:AB_374040

Type: Antibody

Proper Citation

GeneTex Cat# GTX10850, RRID:AB_374040

Antibody Information

URL: http://antibodyregistry.org/AB_374040

Proper Citation: GeneTex Cat# GTX10850, RRID:AB_374040

Target Antigen: TGF beta 2 antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG ELISA, Immunohistochemistry, Neutralizing, Western blot. ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit for TGF beta 2 and TGF beta 1.2 is approximately 0.6 ng/well. IHC: Use at a concentration of 15 µg/ml for paraffin-embedded human tissue sections. Neut: Use at a concentration of 0.01 - 0.03 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit for TGF beta 2 and TGF beta 1.2 is approximately 5 ng/lane and 25 ng/lane under non-reducing and reducing conditions, respectively. Predicted molecular weight: 104 kDa. Anti-TGF beta 2 neutralizes the bioactivity of TGF beta 2 by inhibiting cell proliferation using the IL4 dependent murine HT-2 cell line. In this bioassay, porcine TGF beta 2 (1.0 ng/ml) is preincubated with variou

Antibody Name: TGF beta 2 antibody

Description: This polyclonal targets TGF beta 2 antibody

Target Organism: porcine, human

Antibody ID: AB_374040

Vendor: GeneTex

Catalog Number: GTX10850

Record Creation Time: 20231110T081236+0000

Ratings and Alerts

No rating or validation information has been found for TGF beta 2 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG ELISA, Immunohistochemistry, Neutralizing, Western blot. ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit for TGF beta 2 and TGF beta 1.2 is approximately 0.6 ng/well. IHC: Use at a concentration of 15 µg/ml for paraffin-embedded human tissue sections. Neut: Use at a concentration of 0.01 - 0.03 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit for TGF beta 2 and TGF beta 1.2 is approximately 5 ng/lane and 25 ng/lane under non-reducing and reducing conditions, respectively. Predicted molecular weight: 104 kDa. Anti-TGF beta 2 neutralizes the bioactivity of TGF beta 2 by inhibiting cell proliferation using the IL4 dependent murine HT-2 cell line. In this bioassay, porcine TGF beta 2 (1.0 ng/ml) is preincubated with variou

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.