

Resource Summary Report

Generated by [RRID](#) on Aug 3, 2026

TGF beta 3 antibody

RRID:AB_374041

Type: Antibody

Proper Citation

GeneTex Cat# GTX10851, RRID:AB_374041

Antibody Information

URL: http://antibodyregistry.org/AB_374041

Proper Citation: GeneTex Cat# GTX10851, RRID:AB_374041

Target Antigen: TGF beta 3 antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; ELISA; Western Blot; ELISA: Use at an assay dependent concentration. The detection limit for TGF beta 3 is approximately 0.6 ng/well. Neut: Use at a concentration of 0.005 - 0.015 µg/ml. WB: Use at an assay dependent concentration. The detection limit for TGF beta 3 is approximately 5 ng/lane under non reducing and reducing conditions. Predicted molecular weight: 47 kDa. Anti TGF beta 3 neutralizes the bioactivity of TGF beta 3 by inhibiting cell proliferation using the IL4 dependent murine HT2 cell line. In this bioassay, recombinant human TGF beta 3 (0.1 ng/ml) is preincubated with various concentrations of the antibody (0.0001 to 10 ng/ml) for 1 hour at room temperature in a 96-well microtiter plate. Following this preincubation, HT2 cells are added to ea

Antibody Name: TGF beta 3 antibody

Description: This polyclonal targets TGF beta 3 antibody

Target Organism: chickenbird, human

Antibody ID: AB_374041

Vendor: GeneTex

Catalog Number: GTX10851

Record Creation Time: 20250820T013127+0000

Record Last Update: 20250820T110929+0000

Ratings and Alerts

No rating or validation information has been found for TGF beta 3 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; ELISA; Western Blot; ELISA: Use at an assay dependent concentration. The detection limit for TGF beta 3 is approximately 0.6 ng/well. Neut: Use at a concentration of 0.005 - 0.015 µg/ml. WB: Use at an assay dependent concentration. The detection limit for TGF beta 3 is approximately 5 ng/lane under non reducing and reducing conditions. Predicted molecular weight: 47 kDa. Anti TGF beta 3 neutralizes the bioactivity of TGF beta 3 by inhibiting cell proliferation using the IL4 dependent murine HT2 cell line. In this bioassay, recombinant human TGF beta 3 (0.1 ng/ml) is preincubated with various concentrations of the antibody (0.0001 to 10 ng/ml) for 1 hour at room temperature in a 96-well microtiter plate. Following this preincubation, HT2 cells are added to ea

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.