

Resource Summary Report

Generated by [RRID](#) on Aug 6, 2026

Salivary Amylase antibody

RRID:AB_374557

Type: Antibody

Proper Citation

GeneTex Cat# GTX28944, RRID:AB_374557

Antibody Information

URL: http://antibodyregistry.org/AB_374557

Proper Citation: GeneTex Cat# GTX28944, RRID:AB_374557

Target Antigen: Salivary Amylase antibody

Host Organism: sheep

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Other; Western Blot; ELISA; Immunodiffusion; For western blotting: Antigen: 50 salivaAntibody: 1:5000Secondary Antibody: 1:1000 Donkey a-Sheep/Goat immunoglobulins-peroxidaseDouble diffusion: 10µL antiserum vs 5µL salivaEP: 100µL antiserum vs 5µL saliva Results can be enhanced by washing completed agarose gels in saline, drying and finally stainingwith a protein stain such as coomassie blue. The use of 3 PEG 6000 with 1.2 agarose in a suitable buffer (such as TBE or Tris-barbital pH >8.2) is recommended. Optimal dilutions should be determined by the individual laboratory Suitability for use in nephelometry, immunohistochemical procedures and enzyme-linked immunosorbent assays has notbeen assessed but use in such assays should not necessarily

Antibody Name: Salivary Amylase antibody

Description: This polyclonal targets Salivary Amylase antibody

Target Organism: human

Antibody ID: AB_374557

Vendor: GeneTex

Catalog Number: GTX28944

Record Creation Time: 20250819T213057+0000

Record Last Update: 20250819T231222+0000

Ratings and Alerts

No rating or validation information has been found for Salivary Amylase antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunohistochemistry; Other; Western Blot; ELISA; Immunodiffusion; For western blotting: Antigen: 50 salivaAntibody: 1:5000Secondary Antibody: 1:1000 Donkey a-Sheep/Goat immunoglobulins-peroxidaseDouble diffusion: 10µL antiserum vs 5µL salivaEP: 100µL antiserum vs 5µL saliva Results can be enhanced by washing completed agarose gels in saline, drying and finally stainingwith a protein stain such as coomassie blue. The use of 3 PEG 6000 with 1.2 agarose in a suitable buffer (such as TBE or Tris-barbital pH >8.2) is recommended. Optimal dilutions should be determined by the individual laboratory Suitability for use in nephelometry, immunohistochemical procedures and enzyme-linked immunosorbent assays has notbeen assessed but use in such assays should not necessarily

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.