

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

Osteoprotegerin antibody

RRID:AB_378048

Type: Antibody

Proper Citation

GeneTex Cat# GTX10962, RRID:AB_378048

Antibody Information

URL: http://antibodyregistry.org/AB_378048

Proper Citation: GeneTex Cat# GTX10962, RRID:AB_378048

Target Antigen: Osteoprotegerin antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; ELISA; Flow Cytometry; Western Blot; ELISA, Neutralizing, Western blot. ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit for mouse osteoprotegerin is approximately 0.1 ng/well. Neut: To measure the ability of the antibody to neutralize the bioactivity of mouse OPG, soluble recombinant mouse OPG/Fc is incubated with various concentrations of the antibody in media containing actinomycin D for 1 hour at 37 °C in 96 well plates. WB: Use at a concentration of 1 - 2 µg/ml. This concentration detects mouse OPG at approximately 2 ng/lane and 5 ng/lane under nonreducing and reducing conditions, respectively. Reduced mouse OPG/Fc has a calculated molecular mass of approximately 70.9 kDa. As a result of glycosylation, the recombinant pro

Antibody Name: Osteoprotegerin antibody

Description: This polyclonal targets Osteoprotegerin antibody

Target Organism: rat, mouse

Antibody ID: AB_378048

Vendor: GeneTex

Catalog Number: GTX10962

Record Creation Time: 20250820T010222+0000

Record Last Update: 20250820T095208+0000

Ratings and Alerts

No rating or validation information has been found for Osteoprotegerin antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Block/Neutralize/Inhibit; ELISA; Flow Cytometry; Western Blot; ELISA, Neutralizing, Western blot. ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit for mouse osteoprotegerin is approximately 0.1 ng/well. Neut: To measure the ability of the antibody to neutralize the bioactivity of mouse OPG, soluble recombinant mouse OPG/Fc is incubated with various concentrations of the antibody in media containing actinomycin D for 1 hour at 37 °C in 96 well plates. WB: Use at a concentration of 1 - 2 µg/ml. This concentration detects mouse OPG at approximately 2 ng/lane and 5 ng/lane under nonreducing and reducing conditions, respectively. Reduced mouse OPG/Fc has a calculated molecular mass of approximately 70.9 kDa. As a result of glycosylation, the recombinant pro

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.