

Resource Summary Report

Generated by [RRID](#) on Sep 10, 2026

FOXA1 / HNF3A antibody

RRID:AB_380471

Type: Antibody

Proper Citation

GeneTex Cat# GTX25089, RRID:AB_380471

Antibody Information

URL: http://antibodyregistry.org/AB_380471

Proper Citation: GeneTex Cat# GTX25089, RRID:AB_380471

Target Antigen: FOXA1 / HNF3A antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Approx 50kDa band observed in Human lysates of cell line HepG2 (calculated MW of 49.1kDa according to NP_004487.2). Recommended concentration: 0.2-0.6µg/ml. An additional band of unknown identity was also consistently observed at 35kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? Immunohistochemistry: In paraffin embedded Human Prostate shows nuclear staining of secretory cells. Recommended concentration, 5-10 µg/ml.

Antibody Name: FOXA1 / HNF3A antibody

Description: This polyclonal targets FOXA1 / HNF3A antibody

Target Organism: rat, porcine, cow, pig, mouse, bovine, human

Antibody ID: AB_380471

Vendor: GeneTex

Catalog Number: GTX25089

Record Creation Time: 20231110T081149+0000

Record Last Update: 20260831T231028+0000

Ratings and Alerts

No rating or validation information has been found for FOXA1 / HNF3A antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Approx 50kDa band observed in Human lysates of cell line HepG2 (calculated MW of 49.1kDa according to NP_004487.2). Recommended concentration: 0.2-0.6µg/ml. An additional band of unknown identity was also consistently observed at 35kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? Immunohistochemistry: In paraffin embedded Human Prostate shows nuclear staining of secretory cells. Recommended concentration, 5-10 µg/ml.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.