

Resource Summary Report

Generated by [RRID](#) on Aug 11, 2026

D-Dimer [DD10] antibody

RRID:AB_380838

Type: Antibody

Proper Citation

GeneTex Cat# GTX10051, RRID:AB_380838

Antibody Information

URL: http://antibodyregistry.org/AB_380838

Proper Citation: GeneTex Cat# GTX10051, RRID:AB_380838

Target Antigen: D-Dimer [DD10] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Western Blot; Immunohistochemistry; ELISA; ELISA, Immunoassay, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IA, WB, ELISA: Use at an assay dependent dilution. IA: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. To be analyzed in sandwich immunoassay, plasma must be diluted with 10 mM Tris/HCl buffer, pH 7.5, containing 1 M NaCl and 0.1 Tween 20. To avoid non-specific binding, the final NaCl concentration in plasma samples must be 0.5 M or more. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: D-Dimer [DD10] antibody

Description: This monoclonal targets D-Dimer [DD10] antibody

Target Organism: mouse, human

Antibody ID: AB_380838

Vendor: GeneTex

Catalog Number: GTX10051

Record Creation Time: 20250820T013913+0000

Record Last Update: 20250820T113006+0000

Ratings and Alerts

No rating or validation information has been found for D-Dimer [DD10] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Western Blot; Immunohistochemistry; ELISA; ELISA, Immunoassay, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IA, WB, ELISA: Use at an assay dependent dilution. IA: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. To be analyzed in sandwich immunoassay, plasma must be diluted with 10 mM Tris/HCl buffer, pH 7.5, containing 1 M NaCl and 0.1 Tween 20. To avoid non-specific binding, the final NaCl concentration in plasma samples must be 0.5 M or more. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.