

Resource Summary Report

Generated by [RRID](#) on Sep 7, 2026

D-Dimer [DD5] antibody

RRID:AB_380842

Type: Antibody

Proper Citation

GeneTex Cat# GTX10055, RRID:AB_380842

Antibody Information

URL: http://antibodyregistry.org/AB_380842

Proper Citation: GeneTex Cat# GTX10055, RRID:AB_380842

Target Antigen: D-Dimer [DD5] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: ELISA; Immunohistochemistry; Western Blot; ELISA: Use at an assay dependent dilution. IA: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Can be paired for ELISA with GTX10050 Mouse monoclonal [DD1] to D-Dimer, GTX10052 Mouse monoclonal [DD2] to D-Dimer and GTX10053 Mouse monoclonal [DD3] to D-Dimer . To be analyzed in sandwich immunoassay, plasma must be diluted with 10 mM Tris/HCl buffer, pH 7.5, containing 1 M NaCl and 0.1 Tween 20. To avoid non-specific binding, the final NaCl concentration in plasma samples must be 0.5 M or more. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., ELISA, Immunoassay, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IA,

Antibody Name: D-Dimer [DD5] antibody

Description: This monoclonal targets D-Dimer [DD5] antibody

Target Organism: mouse, human

Antibody ID: AB_380842

Vendor: GeneTex

Catalog Number: GTX10055

Record Creation Time: 20231110T081148+0000

Record Last Update: 20260829T201043+0000

Ratings and Alerts

No rating or validation information has been found for D-Dimer [DD5] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: ELISA; Immunohistochemistry; Western Blot; ELISA: Use at an assay dependent dilution. IA: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Can be paired for ELISA with GTX10050 Mouse monoclonal [DD1] to D-Dimer, GTX10052 Mouse monoclonal [DD2] to D-Dimer and GTX10053 Mouse monoclonal [DD3] to D-Dimer . To be analyzed in sandwich immunoassay, plasma must be diluted with 10 mM Tris/HCl buffer, pH 7.5, containing 1 M NaCl and 0.1 Tween 20. To avoid non-specific binding, the final NaCl concentration in plasma samples must be 0.5 M or more. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user., ELISA, Immunoassay, Western blot. The usefulness of this product in other applications has not been determined., ELISA, IA,

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.