

Resource Summary Report

Generated by [RRID](#) on Sep 6, 2026

CCL4 antibody

RRID:AB_381022

Type: Antibody

Proper Citation

GeneTex Cat# GTX10383, RRID:AB_381022

Antibody Information

URL: http://antibodyregistry.org/AB_381022

Proper Citation: GeneTex Cat# GTX10383, RRID:AB_381022

Target Antigen: CCL4 antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG ELISA; Immunohistochemistry; Block/Neutralize/Inhibit; Western Blot; ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit is approximately 0.16 ng/well. IHC: Use at a concentration of 0.5 - 5 µg/ml to detect MIP-1beta in cultured cells or tissue sections. Neut: Use at a concentration of 0.5 - 2 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit is approximately 1 ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 11 kDa. Neutralization Dose (ND50) is defined as the concentration of antibody required to yield one-half maximal inhibition of the cytokine activity on a responsive cell line, when that cytokine is present at a concentration just high enough to elicit a maximum response. The ND50 is approxi

Antibody Name: CCL4 antibody

Description: This polyclonal targets CCL4 antibody

Target Organism: human

Antibody ID: AB_381022

Vendor: GeneTex

Catalog Number: GTX10383

Record Creation Time: 20231110T080834+0000

Ratings and Alerts

No rating or validation information has been found for CCL4 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG ELISA; Immunohistochemistry; Block/Neutralize/Inhibit; Western Blot; ELISA: Use at a concentration of 0.5 - 1 µg/ml. The detection limit is approximately 0.16 ng/well. IHC: Use at a concentration of 0.5 - 5 µg/ml to detect MIP-1beta in cultured cells or tissue sections. Neut: Use at a concentration of 0.5 - 2 µg/ml. WB: Use at a concentration of 0.1 - 0.2 µg/ml. The detection limit is approximately 1 ng/lane under non-reducing and reducing conditions. Predicted molecular weight: 11 kDa. Neutralization Dose (ND50) is defined as the concentration of antibody required to yield one-half maximal inhibition of the cytokine activity on a responsive cell line, when that cytokine is present at a concentration just high enough to elicit a maximum response. The ND50 is approxi

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.