

Resource Summary Report

Generated by [RRID](#) on Aug 4, 2026

Vimentin antibody

RRID:AB_381287

Type: Antibody

Proper Citation

GeneTex Cat# GTX11256, RRID:AB_381287

Antibody Information

URL: http://antibodyregistry.org/AB_381287

Proper Citation: GeneTex Cat# GTX11256, RRID:AB_381287

Target Antigen: Vimentin antibody

Host Organism: goat

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG ICC: Use at an assay dependent dilution on normal and pathological tissue of mesenchymal derivation. IF: Use at a dilution of 1/20. Can be used with immunoperoxidase or immunofluorescent labeling on formalin-fixed, paraffin-embedded as well as frozen tissue sections to localizes vimentin in fibroblasts, endothelial cells, lymphoid tissues and melanocytes, as well as tumors derived from these cells including sarcomas, lymphomas, melanomas, and their metastatic lesions. WB: Use at an assay dependent dilution. The antibody specifically stains the vimentin band using total cell or tissue extracts. Detects a band of approximately 58 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunocytochemistry, Immunofluorescence, Western blot

Antibody Name: Vimentin antibody

Description: This polyclonal targets Vimentin antibody

Target Organism: feline, rat, hamster, donkey, porcine, canine, goat, horse, mouse, rabbit, bovine, human, sheep

Antibody ID: AB_381287

Vendor: GeneTex

Catalog Number: GTX11256

Record Creation Time: 20250819T230732+0000

Record Last Update: 20250820T042152+0000

Ratings and Alerts

No rating or validation information has been found for Vimentin antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG ICC: Use at an assay dependent dilution on normal and pathological tissue of mesenchymal derivation. IF: Use at a dilution of 1/20. Can be used with immunoperoxidase or immunofluorescent labeling on formalin-fixed, paraffin-embedded as well as frozen tissue sections to localizes vimentin in fibroblasts, endothelial cells, lymphoid tissues and melanocytes, as well as tumors derived from these cells including sarcomas, lymphomas, melanomas, and their metastatic lesions. WB: Use at an assay dependent dilution. The antibody specifically stains the vimentin band using total cell or tissue extracts. Detects a band of approximately 58 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Immunocytochemistry, Immunofluorescence, Western blot

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.