

# Resource Summary Report

Generated by [RRID](#) on Aug 30, 2026

## GJA1 antibody

RRID:AB\_381300

Type: Antibody

---

### Proper Citation

GeneTex Cat# GTX11370, RRID:AB\_381300

---

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_381300](http://antibodyregistry.org/AB_381300)

**Proper Citation:** GeneTex Cat# GTX11370, RRID:AB\_381300

**Target Antigen:** GJA1 antibody

**Host Organism:** rabbit

**Clonality:** polyclonal

**Comments:** Discontinued; manufacturer recommendations: IgG; IgG Immunocytochemistry, Immunofluorescence, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot. ICC: Use at an assay dependent dilution. IF: Use at a dilution of 1/400 (indirect immunofluorescent staining of acetone-fixedcultured baby hamster kidney (BHK)) - 1/2000 (indirect immunofluorescent staining of rat heart (Negative on rat liver sections)). IHC-P: Use at a dilution of 1/2000. Perform enzymatic antigen retrieval before commencing with IHC staining protocol. IHC-Fr: Use at an assay dependent dilution. WB: Use at a dilution of 1/8000. Detects a band of approximately 43 kDa, plus 2-3 smaller weak bands. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.; Western Blot; Immunocytochemistry; Immunohist

**Antibody Name:** GJA1 antibody

**Description:** This polyclonal targets GJA1 antibody

**Target Organism:** rat, hamster, cow, mouse, chickenbird, bovine, human

**Antibody ID:** AB\_381300

**Vendor:** GeneTex

**Catalog Number:** GTX11370

**Record Creation Time:** 20250819T224659+0000

**Record Last Update:** 20250820T031723+0000

---

## Ratings and Alerts

No rating or validation information has been found for GJA1 antibody.

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunocytochemistry, Immunofluorescence, IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections), Western blot. ICC: Use at an assay dependent dilution. IF: Use at a dilution of 1/400 (indirect immunofluorescent staining of acetone-fixedcultured baby hamster kidney (BHK)) - 1/2000 (indirect immunofluorescent staining of rat heart (Negative on rat liver sections)). IHC-P: Use at a dilution of 1/2000. Perform enzymatic antigen retrieval before commencing with IHC staining protocol. IHC-Fr: Use at an assay dependent dilution. WB: Use at a dilution of 1/8000. Detects a band of approximately 43 kDa, plus 2-3 smaller weak bands. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.; Western Blot; Immunocytochemistry; Immunohist

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We have not found any literature mentions for this resource.