

Resource Summary Report

Generated by [RRID](#) on Aug 7, 2026

HDAC7 antibody

RRID:AB_381381

Type: Antibody

Proper Citation

GeneTex Cat# GTX12175, RRID:AB_381381

Antibody Information

URL: http://antibodyregistry.org/AB_381381

Proper Citation: GeneTex Cat# GTX12175, RRID:AB_381381

Target Antigen: HDAC7 antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunoprecipitation, Western blot. IP: Use at an assay dependent dilution. 1.5-1.8 µg of the antibody immunoprecipitates HDAC7 from a RIPA extract (180 µg) of 293T human embryonic kidney cells expressing recombinant human HDAC7. WB: Use at an assay dependent dilution. Predicted molecular weight: 105 kDa. Additional weak bands may be detected in various extract preparations. Detection of HDAC7 is specifically inhibited with the immunizing peptide. A minimum working antibody dilution of 1/2000 is recommended using whole extracts of mouse NIH-3T3 cells. A minimum working antibody dilution of 1/500 is recommended using whole extracts of rat PC12 cells. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.; Western Blot; Immunopr

Antibody Name: HDAC7 antibody

Description: This polyclonal targets HDAC7 antibody

Target Organism: rat, mouse, human

Antibody ID: AB_381381

Vendor: GeneTex

Catalog Number: GTX12175

Record Creation Time: 20250819T223440+0000

Record Last Update: 20250820T023809+0000

Ratings and Alerts

No rating or validation information has been found for HDAC7 antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunoprecipitation, Western blot. IP: Use at an assay dependent dilution. 1.5-1.8 µg of the antibody immunoprecipitates HDAC7 from a RIPA extract (180 µg) of 293T human embryonic kidney cells expressing recombinant human HDAC7. WB: Use at an assay dependent dilution. Predicted molecular weight: 105 kDa. Additional weak bands may be detected in various extract preparations. Detection of HDAC7 is specifically inhibited with the immunizing peptide. A minimum working antibody dilution of 1/2000 is recommended using whole extracts of mouse NIH-3T3 cells. A minimum working antibody dilution of 1/500 is recommended using whole extracts of rat PC12 cells. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.; Western Blot; Immunopr

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.