

Resource Summary Report

Generated by [RRID](#) on Aug 7, 2026

Cytohesin 1 antibody [CYT1-82]

RRID:AB_381425

Type: Antibody

Proper Citation

GeneTex Cat# GTX12425, RRID:AB_381425

Antibody Information

URL: http://antibodyregistry.org/AB_381425

Proper Citation: GeneTex Cat# GTX12425, RRID:AB_381425

Target Antigen: Cytohesin 1 antibody [CYT1-82]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG1; IgG1 Western Blot; ELISA; Immunocytochemistry; ELISA, Immunocytochemistry, Western blot. The usefulness of this product in other applications has not been determined., ELISA, ICC, WB, ELISA: Use at an assay dependent dilution. ICC: Use at an assay dependent dilution. WB: Use at a concentration of 2 - 4 µg/ml. This concentration is determined by using a whole extract of cultured human lymphoma Raji cells. A band is detected at approximately 50 kDa, and also additional bands at approximately 30 and 90 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Antibody Name: Cytohesin 1 antibody [CYT1-82]

Description: This monoclonal targets Cytohesin 1 antibody [CYT1-82]

Target Organism: human

Antibody ID: AB_381425

Vendor: GeneTex

Catalog Number: GTX12425

Record Creation Time: 20250820T035808+0000

Record Last Update: 20250820T174043+0000

Ratings and Alerts

No rating or validation information has been found for Cytohesin 1 antibody [CYT1-82].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG1; IgG1 Western Blot; ELISA; Immunocytochemistry; ELISA, Immunocytochemistry, Western blot. The usefulness of this product in other applications has not been determined., ELISA, ICC, WB, ELISA: Use at an assay dependent dilution. ICC: Use at an assay dependent dilution. WB: Use at a concentration of 2 - 4 µg/ml. This concentration is determined by using a whole extract of cultured human lymphoma Raji cells. A band is detected at approximately 50 kDa, and also additional bands at approximately 30 and 90 kDa. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.