

Resource Summary Report

Generated by [RRID](#) on Aug 10, 2026

Bcl-X [2H12] antibody

RRID:AB_383754

Type: Antibody

Proper Citation

GeneTex Cat# GTX73931, RRID:AB_383754

Antibody Information

URL: http://antibodyregistry.org/AB_383754

Proper Citation: GeneTex Cat# GTX73931, RRID:AB_383754

Target Antigen: Bcl-X [2H12] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Immunohistochemistry; Western Blot; Immunoprecipitation; Immunohistochemistry - fixed; This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining. Can also be used for: Immunoprecipitation (1:50), Western Blotting (1:50 to 1:100), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Antibody Name: Bcl-X [2H12] antibody

Description: This monoclonal targets Bcl-X [2H12] antibody

Antibody ID: AB_383754

Vendor: GeneTex

Catalog Number: GTX73931

Record Creation Time: 20250820T012300+0000

Record Last Update: 20250820T104658+0000

Ratings and Alerts

No rating or validation information has been found for Bcl-X [2H12] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry; Western Blot; Immunoprecipitation; Immunohistochemistry - fixed; This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining. Can also be used for: Immunoprecipitation (1:50), Western Blotting (1:50 to 1:100), IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.