

# Resource Summary Report

Generated by [RRID](#) on Sep 7, 2026

## MVP [1032] antibody

RRID:AB\_383812

Type: Antibody

### Proper Citation

GeneTex Cat# GTX74004, RRID:AB\_383812

### Antibody Information

**URL:** [http://antibodyregistry.org/AB\\_383812](http://antibodyregistry.org/AB_383812)

**Proper Citation:** GeneTex Cat# GTX74004, RRID:AB\_383812

**Target Antigen:** MVP [1032] antibody

**Host Organism:** mouse

**Clonality:** monoclonal

**Comments:** Discontinued; manufacturer recommendations: Immunohistochemistry - fixed; Immunohistochemistry; Immunoprecipitation; Western Blot; IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC-P, IP, WB, This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the end user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to i

**Antibody Name:** MVP [1032] antibody

**Description:** This monoclonal targets MVP [1032] antibody

**Antibody ID:** AB\_383812

**Vendor:** GeneTex

**Catalog Number:** GTX74004

**Record Creation Time:** 20231110T081209+0000

**Record Last Update:** 20260901T000132+0000

---

## Ratings and Alerts

No rating or validation information has been found for MVP [1032] antibody.

**Warning:** Discontinued at GeneTex

Discontinued; manufacturer recommendations: Immunohistochemistry - fixed; Immunohistochemistry; Immunoprecipitation; Western Blot; IHC (Formalin-fixed paraffin-embedded sections), Immunoprecipitation, Western blot. The usefulness of this product in other applications has not been determined., IHC-P, IP, WB, This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the end user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to i

---

## Data and Source Information

**Source:** [Antibody Registry](#)

---

## Usage and Citation Metrics

We have not found any literature mentions for this resource.