

Resource Summary Report

Generated by [RRID](#) on Aug 2, 2026

Blood Group Lewis y [F3] antibody

RRID:AB_384071

Type: Antibody

Proper Citation

GeneTex Cat# GTX23359, RRID:AB_384071

Antibody Information

URL: http://antibodyregistry.org/AB_384071

Proper Citation: GeneTex Cat# GTX23359, RRID:AB_384071

Target Antigen: Blood Group Lewis y [F3] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Ig3 Flow Cytometry; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; Immunofluorescence; Flow cytometry/FACS, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., FACS, IHC-P, WB, For FACS: Use at an assay dependent dilution. For IHC-P: Use at a dilution of 1:40 or greater, for Biotin-based detection systems. For optimal staining, the primary antibody should be incubated 20-60 minutes at RT. Pre-treatment not required. For WB: Use at an assay dependent dilution. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher.

Antibody Name: Blood Group Lewis y [F3] antibody

Description: This monoclonal targets Blood Group Lewis y [F3] antibody

Target Organism: human

Antibody ID: AB_384071

Vendor: GeneTex

Catalog Number: GTX23359

Record Creation Time: 20250820T040053+0000

Record Last Update: 20250820T174814+0000

Ratings and Alerts

No rating or validation information has been found for Blood Group Lewis y [F3] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Ig3 Flow Cytometry; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; Immunofluorescence; Flow cytometry/FACS, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications has not been determined., FACS, IHC-P, WB, For FACS: Use at an assay dependent dilution. For IHC-P: Use at a dilution of 1:40 or greater, for Biotin-based detection systems. For optimal staining, the primary antibody should be incubated 20-60 minutes at RT. Pre-treatment not required. For WB: Use at an assay dependent dilution. Not tested in other applications. Optimal dilutions/concentrations should be determined by the researcher.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.