

Resource Summary Report

Generated by [RRID](#) on Sep 10, 2026

Progesterone antibody

RRID:AB_384200

Type: Antibody

Proper Citation

GeneTex Cat# GTX11002, RRID:AB_384200

Antibody Information

URL: http://antibodyregistry.org/AB_384200

Proper Citation: GeneTex Cat# GTX11002, RRID:AB_384200

Target Antigen: Progesterone antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG RIA: The antiserum is characterized utilizing the following dextran coated charcoal radioimmunoassay (RIA) protocol, where 0.5 ml of reconstituted and diluted antiserum has been found to bind at least 40 of 10-15 picograms of tritiated (³H) progesterone with a specific activity of approximately 100 Ci/mmole. RIA Reagents(A) Standards: Prepare a stock standard solution of 1 µg/ml progesterone in absolute ethanol. Dilute a portion of the stock solution with buffer (B) to a concentration of 500 pg/0.1ml. This is further diluted in buffer (B) to obtain standard solutions at the following concentrations: 250, 125, 63, 31 and 15 pg/0.1 ml.(B) Dilution buffer: 0.05 M Tris-HCl, pH 8.0, containing 0.1 M NaCl, 0.1 BSA and 0.1 sodium azide.(C) Dextran coated charcoal suspension: 0.5 activa

Antibody Name: Progesterone antibody

Description: This polyclonal targets Progesterone antibody

Target Organism: human

Antibody ID: AB_384200

Vendor: GeneTex

Catalog Number: GTX11002

Record Creation Time: 20231110T080837+0000

Ratings and Alerts

No rating or validation information has been found for Progesterone antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG RIA: The antiserum is characterized utilizing the following dextran coated charcoal radioimmunoassay (RIA) protocol, where 0.5 ml of reconstituted and diluted antiserum has been found to bind at least 40 of 10-15 picograms of tritiated (³H) progesterone with a specific activity of approximately 100 Ci/mmol. RIA Reagents(A) Standards: Prepare a stock standard solution of 1 µg/ml progesterone in absolute ethanol. Dilute a portion of the stock solution with buffer (B) to a concentration of 500 pg/0.1ml. This is further diluted in buffer (B) to obtain standard solutions at the following concentrations: 250, 125, 63, 31 and 15 pg/0.1 ml.(B) Dilution buffer: 0.05 M Tris-HCl, pH 8.0, containing 0.1 M NaCl, 0.1 BSA and 0.1 sodium azide.(C) Dextran coated charcoal suspension: 0.5 activa

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.