

Resource Summary Report

Generated by [RRID](#) on Jul 31, 2026

Triadin antibody [GE 4.90]

RRID:AB_384889

Type: Antibody

Proper Citation

GeneTex Cat# GTX22870, RRID:AB_384889

Antibody Information

URL: http://antibodyregistry.org/AB_384889

Proper Citation: GeneTex Cat# GTX22870, RRID:AB_384889

Target Antigen: Triadin antibody [GE 4.90]

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG For IHC: use at a dilution of 1/50. For WB: use at a dilution of 1/500. For IP: Assay dependent. ELISA: Assay dependent. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects a 95 kDa protein representing Triadin from rabbit skeletal muscle extracts. Immunohistochemical staining of Triadin in rat skeletal muscle results in striated staining consistent with sarcoplasmic reticulum localization., ELISA, IHC, IHC-Fr, IP, WB, ELISA, Immunohistochemistry, IHC (Frozen sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined.; Western Blot; Immunohistochemistry - frozen; Immunohistochemistry; ELISA; Immunoprecipitation

Antibody Name: Triadin antibody [GE 4.90]

Description: This monoclonal targets Triadin antibody [GE 4.90]

Target Organism: rat, mouse, rabbit

Antibody ID: AB_384889

Vendor: GeneTex

Catalog Number: GTX22870

Record Creation Time: 20250820T024602+0000

Ratings and Alerts

No rating or validation information has been found for Triadin antibody [GE 4.90].

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG For IHC: use at a dilution of 1/50. For WB: use at a dilution of 1/500. For IP: Assay dependent. ELISA: Assay dependent. Not tested in other applications. Optimal dilutions / concentrations should be determined by the end user. By Western blot, this antibody detects a 95 kDa protein representing Triadin from rabbit skeletal muscle extracts. Immunohistochemical staining of Triadin in rat skeletal muscle results in striated staining consistent with sarcoplasmic reticulum localization., ELISA, IHC, IHC-Fr, IP, WB, ELISA, Immunohistochemistry, IHC (Frozen sections), Immunoprecipitation, Western Blot. The usefulness of this product in other applications has not been determined.; Western Blot; Immunohistochemistry - frozen; Immunohistochemistry; ELISA; Immunoprecipitation

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.