

Resource Summary Report

Generated by [RRID](#) on Aug 9, 2026

Cytokeratin 10 + 13 [DE-K13] antibody

RRID:AB_385885

Type: Antibody

Proper Citation

GeneTex Cat# GTX29383, RRID:AB_385885

Antibody Information

URL: http://antibodyregistry.org/AB_385885

Proper Citation: GeneTex Cat# GTX29383, RRID:AB_385885

Target Antigen: Cytokeratin 10 + 13 [DE-K13] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Flow Cytometry; Western Blot; Immunohistochemistry - frozen; Immunocytochemistry; Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry; FACS: 1/100 - 1/200. ICC: Use at an assay dependent dilution. IHC-Fr: Use at an assay dependent dilution. IHC-P: Use at an assay dependent dilution. DE-K13 reacts with cytokeratin 10 and 13 in fixed cells and on frozen tissues, but exclusively with cytokeratin 13 in formalin-fixed, paraffin-embedded tissues. For formalin fixed paraffin embedded tissue, pretreatment with 0.1 pepsin in 0.1M HCL for 30 minutes at room temperature is required. Recommended range is 1:100 - 1:200 for immunohistochemistry with avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent. WB: 1/100 - 1/1000., Flow cytometry/FACS, Immunocy

Antibody Name: Cytokeratin 10 + 13 [DE-K13] antibody

Description: This monoclonal targets Cytokeratin 10 + 13 [DE-K13] antibody

Target Organism: feline, human

Antibody ID: AB_385885

Vendor: GeneTex

Catalog Number: GTX29383

Record Creation Time: 20250820T060523+0000

Ratings and Alerts

No rating or validation information has been found for Cytokeratin 10 + 13 [DE-K13] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Flow Cytometry; Western Blot; Immunohistochemistry - frozen; Immunocytochemistry; Immunohistochemistry - fixed; Immunofluorescence; Immunohistochemistry; FACS: 1/100 - 1/200. ICC: Use at an assay dependent dilution. IHC-Fr: Use at an assay dependent dilution. IHC-P: Use at an assay dependent dilution. DE-K13 reacts with cytokeratin 10 and 13 in fixed cells and on frozen tissues, but exclusively with cytokeratin 13 in formalin-fixed, paraffin-embedded tissues. For formalin fixed paraffin embedded tissue, pretreatment with 0.1 pepsin in 0.1M HCL for 30 minutes at room temperature is required. Recommended range is 1:100 - 1:200 for immunohistochemistry with avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent. WB: 1/100 - 1/1000., Flow cytometry/FACS, Immunocy

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.