

Resource Summary Report

Generated by [RRID](#) on Sep 11, 2026

NCAM [1213C3.D5] antibody (ready-to-use)

RRID:AB_422643

Type: Antibody

Proper Citation

GeneTex Cat# GTX17235, RRID:AB_422643

Antibody Information

URL: http://antibodyregistry.org/AB_422643

Proper Citation: GeneTex Cat# GTX17235, RRID:AB_422643

Target Antigen: NCAM [1213C3.D5] antibody (ready-to-use)

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: IHC-F, IHC-Fr, IHC-P, IHC (Formalin fixed sections), IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-P: Pre-diluted for an incubation period of 30 minutes at room temperature. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. IHC-Fr: Pre-diluted for an incubation period of 30 minutes at room temperature. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry

Antibody Name: NCAM [1213C3.D5] antibody (ready-to-use)

Description: This monoclonal targets NCAM [1213C3.D5] antibody (ready-to-use)

Target Organism: human

Antibody ID: AB_422643

Vendor: GeneTex

Catalog Number: GTX17235

Record Creation Time: 20231110T081103+0000

Record Last Update: 20260831T204936+0000

Ratings and Alerts

No rating or validation information has been found for NCAM [1213C3.D5] antibody (ready-to-use).

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IHC-F, IHC-Fr, IHC-P, IHC (Formalin fixed sections), IHC (Frozen sections), IHC (Formalin-fixed paraffin-embedded sections). The usefulness of this product in other applications has not been determined., IHC-P: Pre-diluted for an incubation period of 30 minutes at room temperature. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. IHC-Fr: Pre-diluted for an incubation period of 30 minutes at room temperature. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.; Immunohistochemistry - frozen; Immunohistochemistry - fixed; Immunohistochemistry

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.