

Resource Summary Report

Generated by [RRID](#) on Sep 7, 2026

MAdCAM1 [17F5] antibody

RRID:AB_625356

Type: Antibody

Proper Citation

GeneTex Cat# GTX54143, RRID:AB_625356

Antibody Information

URL: http://antibodyregistry.org/AB_625356

Proper Citation: GeneTex Cat# GTX54143, RRID:AB_625356

Target Antigen: MAdCAM1 [17F5] antibody

Host Organism: mouse

Clonality: monoclonal

Comments: Discontinued; manufacturer recommendations: Flow Cytometry; ELISA; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; For ELISA: Use at an assay dependent dilution. For FACS: Use 10ul of the suggested working dilution (1:10-1:100) to label 10^6 cells in 100ul. For IHC-P: Use at 1:50-1:100. This product does not require protein digestion prior to staining. This product does not require HIER prior to staining. For WB: Use at an assay dependent dilution under reducing conditions. Detects a band of approximately 60 kDa (Predicted molecular weight: 42 kDa). Not yet tested in other applications. Optimal dilutions/concentrations should be determined by the end user., ELISA, Flow cytometry/FACS, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications

Antibody Name: MAdCAM1 [17F5] antibody

Description: This monoclonal targets MAdCAM1 [17F5] antibody

Target Organism: human

Antibody ID: AB_625356

Vendor: GeneTex

Catalog Number: GTX54143

Record Creation Time: 20231110T080404+0000

Ratings and Alerts

No rating or validation information has been found for MAdCAM1 [17F5] antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: Flow Cytometry; ELISA; Immunofluorescence; Immunohistochemistry; Immunohistochemistry - fixed; Western Blot; For ELISA: Use at an assay dependent dilution. For FACS: Use 10ul of the suggested working dilution (1:10-1:100) to label 10^6 cells in 100ul. For IHC-P: Use at 1:50-1:100. This product does not require protein digestion prior to staining. This product does not require HIER prior to staining. For WB: Use at an assay dependent dilution under reducing conditions. Detects a band of approximately 60 kDa (Predicted molecular weight: 42 kDa). Not yet tested in other applications. Optimal dilutions/concentrations should be determined by the end user., ELISA, Flow cytometry/FACS, IHC (Formalin-fixed paraffin-embedded sections), Western blot. The usefulness of this product in other applications

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.