

Resource Summary Report

Generated by [RRID](#) on Sep 10, 2026

Numb antibody

RRID:AB_626001

Type: Antibody

Proper Citation

GeneTex Cat# GTX30689, RRID:AB_626001

Antibody Information

URL: http://antibodyregistry.org/AB_626001

Proper Citation: GeneTex Cat# GTX30689, RRID:AB_626001

Target Antigen: Numb antibody

Host Organism: rabbit

Clonality: polyclonal

Comments: Discontinued; manufacturer recommendations: IgG; IgG Immunocytochemistry; Western Blot; By Western blot one may see any or all of the four isoforms. The molecular weight of human and mouse numb is 72 kDa (isoform 1), 66 kDa (isoform 2), 71 kDa (isoform 3), and 65 kDa (isoform 4). Multiple non-specific bands may be seen with lower dilutions and/or longer exposure times than suggested in the protocol we used to obtain our data. No other applications have been tested. Recommended Starting Dilutions:* Western Blot: 0.2-0.5 ug/ml Immunoprecipitation: Not tested Immunohisto/cytochemistry: Not tested *Optimal working dilutions should be determined by the investigator. Western Blot.

Antibody Name: Numb antibody

Description: This polyclonal targets Numb antibody

Target Organism: chicken/bird, mouse, human and chicken (isoforms 1-4) proteins and 90 homology with rat numb protein (isoforms unknown), human, human protein. the immunogen used for this antibody production has 100 homology with mouse

Antibody ID: AB_626001

Vendor: GeneTex

Catalog Number: GTX30689

Record Creation Time: 20231110T080427+0000

Record Last Update: 20260901T061140+0000

Ratings and Alerts

No rating or validation information has been found for Numb antibody.

Warning: Discontinued at GeneTex

Discontinued; manufacturer recommendations: IgG; IgG Immunocytochemistry; Western Blot; By Western blot one may see any or all of the four isoforms. The molecular weight of human and mouse numb is 72 kDa (isoform 1), 66 kDa (isoform 2), 71 kDa (isoform 3), and 65 kDa (isoform 4). Multiple non-specific bands may be seen with lower dilutions and/or longer exposure times than suggested in the protocol we used to obtain our data. No other applications have been tested. Recommended Starting Dilutions:* Western Blot: 0.2-0.5 ug/ml Immunoprecipitation: Not tested Immunohisto/cytochemistry: Not tested
*Optimal working dilutions should be determined by the investigator. Western Blot.

Data and Source Information

Source: [Antibody Registry](#)

Usage and Citation Metrics

We have not found any literature mentions for this resource.