

Resource Summary Report

Generated by [RRID](#) on Aug 27, 2026

hnRNPA2_LC_WT

RRID:Addgene_98657

Type: Plasmid

Proper Citation

RRID:Addgene_98657

Plasmid Information

URL: <http://www.addgene.org/98657>

Proper Citation: RRID:Addgene_98657

Insert Name: hnRNPA2

Organism: Homo sapiens

Bacterial Resistance: Kanamycin

Defining Citation: [PMID:29358076](#)

Vector Backbone Description: Vector Backbone:pJ411; Vector Types:Bacterial Expression; Bacterial Resistance:Kanamycin

Plasmid Name: hnRNPA2_LC_WT

Record Creation Time: 20220422T222631+0000

Record Last Update: 20260815T082240+0000

Ratings and Alerts

No rating or validation information has been found for hnRNPA2_LC_WT.

No alerts have been found for hnRNPA2_LC_WT.

Data and Source Information

Source: [Addgene](#)

Usage and Citation Metrics

We found 4 mentions in open access literature.

Listed below are recent publications. The full list is available at [RRID](#) .

Van Lindt J, et al. (2022) F/YGG-motif is an intrinsically disordered nucleic-acid binding motif. *RNA biology*, 19(1), 622.

Perdikari TM, et al. (2020) SARS-CoV-2 nucleocapsid protein undergoes liquid-liquid phase separation stimulated by RNA and partitions into phases of human ribonucleoproteins. *bioRxiv : the preprint server for biology*.

Ryan VH, et al. (2020) Weak binding to the A2RE RNA rigidifies hnRNPA2 RRM and reduces liquid-liquid phase separation and aggregation. *Nucleic acids research*, 48(18), 10542.

Perdikari TM, et al. (2020) SARS-CoV-2 nucleocapsid protein phase-separates with RNA and with human hnRNPs. *The EMBO journal*, 39(24), e106478.