

# Resource Summary Report

Generated by [RRID](#) on Aug 19, 2026

## [pJJR82](#)

RRID:Addgene\_75027

Type: Plasmid

### Proper Citation

RRID:Addgene\_75027

### Plasmid Information

**URL:** <http://www.addgene.org/75027>

**Proper Citation:** RRID:Addgene\_75027

**Insert Name:** EGFP^SEC^3xFlag

**Organism:** Caenorhabditis elegans

**Bacterial Resistance:** Ampicillin

**Defining Citation:** [PMID:32586975](#)

**Vector Backbone Description:** Backbone Size:3854; Vector Backbone:pDD282; Vector Types:Worm Expression, CRISPR; Bacterial Resistance:Ampicillin

**Plasmid Name:** pJJR82

**Record Creation Time:** 20220422T222456+0000

**Record Last Update:** 20260815T081942+0000

### Ratings and Alerts

No rating or validation information has been found for pJJR82.

No alerts have been found for pJJR82.

### Data and Source Information

**Source:** [Addgene](#)

### Usage and Citation Metrics

We found 6 mentions in open access literature.

**Listed below are recent publications.** The full list is available at [RRID](#) .

Cornes E, et al. (2022) piRNAs initiate transcriptional silencing of spermatogenic genes during *C. elegans* germline development. *Developmental cell*, 57(2), 180.

Sepers JJ, et al. (2022) The mIAA7 degron improves auxin-mediated degradation in *Caenorhabditiselegans*. *G3 (Bethesda, Md.)*, 12(10).

Remmelzwaal S, et al. (2021) BBLN-1 is essential for intermediate filament organization and apical membrane morphology. *Current biology : CB*, 31(11), 2334.

Castiglioni VG, et al. (2020) Epidermal PAR-6 and PKC-3 are essential for larval development of *C. elegans* and organize non-centrosomal microtubules. *eLife*, 9.

Vicencio J, et al. (2019) Efficient Generation of Endogenous Fluorescent Reporters by Nested CRISPR in *Caenorhabditis elegans*. *Genetics*, 211(4), 1143.

Lloret-Fernández C, et al. (2018) A transcription factor collective defines the HSN serotonergic neuron regulatory landscape. *eLife*, 7.