

Resource Summary Report

Generated by [RRID](#) on Aug 26, 2026

pDonor-tBFP-NLS-Neo (Universal)

RRID:Addgene_80767

Type: Plasmid

Proper Citation

RRID:Addgene_80767

Plasmid Information

URL: <http://www.addgene.org/80767>

Proper Citation: RRID:Addgene_80767

Insert Name: PITCh-gRNA#3 targeting sequence (GCATCGTACGCGTACGTGTT)

Organism: Synthetic

Bacterial Resistance: Kanamycin

Defining Citation: [PMID:28179459](#)

Vector Backbone Description: Backbone Marker:Kazuhisa Nakayama Lab. (Addgene plasmid #80766); Backbone Size:4761; Vector Backbone:pDonor-tBFP-NLS-Neo; Vector Types:Mammalian Expression, CRISPR; Bacterial Resistance:Kanamycin

Plasmid Name: pDonor-tBFP-NLS-Neo (Universal)

Record Creation Time: 20220422T222527+0000

Record Last Update: 20260815T082041+0000

Ratings and Alerts

No rating or validation information has been found for pDonor-tBFP-NLS-Neo (Universal).

No alerts have been found for pDonor-tBFP-NLS-Neo (Universal).

Data and Source Information

Source: [Addgene](#)

Usage and Citation Metrics

We found 19 mentions in open access literature.

Listed below are recent publications. The full list is available at [RRID](#) .

Liu AZ, et al. (2025) The scaffold protein PRR14L links the PP2A-TACC3 axis to mitotic fidelity and sensitivity to MPS1 inhibition. *bioRxiv : the preprint server for biology*.

Accogli A, et al. (2024) Variants in the WDR44 WD40-repeat domain cause a spectrum of ciliopathy by impairing ciliogenesis initiation. *Nature communications*, 15(1), 365.

Ninomiya K, et al. (2024) Calcium influx promotes PLEKHG4B localization to cell-cell junctions and regulates the integrity of junctional actin filaments. *Molecular biology of the cell*, 35(2), ar24.

Hiyamizu S, et al. (2023) Dynein-2-driven intraciliary retrograde trafficking indirectly requires multiple interactions of IFT54 in the IFT-B complex with the dynein-2 complex. *Biology open*, 12(7).

Takasugi M, et al. (2023) CD44 correlates with longevity and enhances basal ATF6 activity and ER stress resistance. *Cell reports*, 42(9), 113130.

Satoda Y, et al. (2022) BROMI/TBC1D32 together with CCRK/CDK20 and FAM149B1/JBTS36 contributes to intraflagellar transport turnaround involving ICK/CILK1. *Molecular biology of the cell*, 33(9), ar79.

Kawamura A, et al. (2022) DYRK2 maintains genome stability via neddylation of cullins in response to DNA damage. *Journal of cell science*, 135(11).

Ishida Y, et al. (2022) Molecular basis underlying the ciliary defects caused by IFT52 variations found in skeletal ciliopathies. *Molecular biology of the cell*, 33(9), ar83.

Qiu H, et al. (2022) Combinations of deletion and missense variations of the dynein-2 DYNC2LI1 subunit found in skeletal ciliopathies cause ciliary defects. *Scientific reports*, 12(1), 31.

Fujisawa S, et al. (2021) ARL3 and ARL13B GTPases participate in distinct steps of INPP5E targeting to the ciliary membrane. *Biology open*, 10(9).

Noguchi T, et al. (2021) CCRK/CDK20 regulates ciliary retrograde protein trafficking via interacting with BROMI/TBC1D32. *PLoS one*, 16(10), e0258497.

Qiu H, et al. (2021) Interaction of INPP5E with ARL13B is essential for its ciliary membrane retention but dispensable for its ciliary entry. *Biology open*, 10(1).

Okazaki M, et al. (2020) Formation of the B9-domain protein complex MKS1-B9D2-B9D1 is essential as a diffusion barrier for ciliary membrane proteins. *Molecular biology of the cell*, 31(20), 2259.

Tsurumi Y, et al. (2019) Interactions of the dynein-2 intermediate chain WDR34 with the light chains are required for ciliary retrograde protein trafficking. *Molecular biology of the cell*, 30(5), 658.

Nozaki S, et al. (2019) Requirement of IFT-B-BBSome complex interaction in export of GPR161 from cilia. *Biology open*, 8(9).

Hamada Y, et al. (2018) Interaction of WDR60 intermediate chain with TCTEX1D2 light chain of the dynein-2 complex is crucial for ciliary protein trafficking. *Molecular biology of the cell*, 29(13), 1628.

Nozaki S, et al. (2018) BBS1 is involved in retrograde trafficking of ciliary GPCRs in the context of the BBSome complex. *PloS one*, 13(3), e0195005.

Takahara M, et al. (2018) Ciliopathy-associated mutations of IFT122 impair ciliary protein trafficking but not ciliogenesis. *Human molecular genetics*, 27(3), 516.

Takei R, et al. (2018) Robust interaction of IFT70 with IFT52-IFT88 in the IFT-B complex is required for ciliogenesis. *Biology open*, 7(5).